

Powers

Multiplication Index Law	When multiplying with the same base (number or letter), add the powers . $a^m \times a^n = a^{m+n}$	$7^5 \times 7^3 = 7^8$ $a^{12} \times a = a^{13}$ $4x^5 \times 2x^8 = 8x^{13}$
Division Index Law	When dividing with the same base (number or letter), subtract the powers . $a^m \div a^n = a^{m-n}$	$15^7 \div 15^4 = 15^3$ $x^9 \div x^2 = x^7$ $20a^{11} \div 5a^3 = 4a^8$
Brackets Index Laws	When raising a power to another power, multiply the powers together. $(a^m)^n = a^{mn}$	$(y^2)^5 = y^{10}$ $(6^3)^4 = 6^{12}$ $(5x^6)^3 = 125x^{18}$
Notable Powers	$p = p^1$ $p^0 = 1$	$99999^0 = 1$
Negative Powers	A negative power performs the reciprocal. $a^{-m} = \frac{1}{a^m}$	$3^{-2} = \frac{1}{3^2} = \frac{1}{9}$
Fractional Powers	The denominator of a fractional power acts as a 'root'. The numerator of a fractional power acts as a normal power. $a^{\frac{m}{n}} = (\sqrt[n]{a})^m$	$27^{\frac{2}{3}} = (\sqrt[3]{27})^2 = 3^2 = 9$ $\left(\frac{25}{16}\right)^{\frac{3}{2}} = \left(\frac{\sqrt{25}}{\sqrt{16}}\right)^3 = \left(\frac{5}{4}\right)^3 = \frac{125}{64}$

Factorising

Factorising Quadratics	When a quadratic expression is in the form $x^2 + bx + c$ find the two numbers that add to give b and multiply to give c .	$x^2 + 7x + 10 = (x + 5)(x + 2)$ (because 5 and 2 add to give 7 and multiply to give 10) $x^2 + 2x - 8 = (x + 4)(x - 2)$ (because +4 and -2 add to give +2 and multiply to give -8)
Difference of Two Squares	An expression of the form $a^2 - b^2$ can be factorised to give $(a + b)(a - b)$	$x^2 - 25 = (x + 5)(x - 5)$ $16x^2 - 81 = (4x + 9)(4x - 9)$

Fractions

Simplifying Fractions	Divide the numerator and denominator by the highest common factor.	$\frac{20}{45} = \frac{4}{9}$
Equivalent Fractions	Fractions which represent the same value .	$\frac{2}{5} = \frac{4}{10} = \frac{20}{50} = \frac{60}{150}$ etc.
Comparing Fractions	To compare fractions, they each need to be rewritten so that they have a common denominator . Ascending means smallest to biggest . Descending means biggest to smallest .	Put in to ascending order : $\frac{3}{4}, \frac{2}{3}, \frac{5}{6}, \frac{1}{2}$. Equivalent: $\frac{9}{12}, \frac{8}{12}, \frac{10}{12}, \frac{6}{12}$ Correct order: $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{5}{6}$
Adding or Subtracting Fractions	Find the LCM of the denominators to find a common denominator. Use equivalent fractions to change each fraction to the common denominator . Then just add or subtract the numerators and keep the denominator the same .	$\frac{2}{3} + \frac{4}{5}$ Multiples of 3: 3, 6, 9, 12, 15 .. Multiples of 5: 5, 10, 15 .. LCM of 3 and 5 = 15 $\frac{2}{3} = \frac{10}{15}$ and $\frac{4}{5} = \frac{12}{15}$ $\frac{10}{15} + \frac{12}{15} = \frac{22}{15} = 1\frac{7}{15}$

Estimating

Error Interval	A range of values that a number could have taken before being rounded or truncated. An error interval is written using inequalities, with a lower bound and an upper bound . Note that the lower bound inequality can be 'equal to', but the upper bound cannot be 'equal to'.	0.6 has been rounded to 1 decimal place. The error interval is: $0.55 \leq x < 0.65$ The lower bound is 0.55 The upper bound is 0.65
Estimate	To find something close to the correct answer .	An estimate for the height of a man is 1.8 metres.
Approximation	When using approximations to estimate the solution to a calculation, round each number in the calculation to 1 significant figure . \approx means 'approximately equal to'	$\frac{348 + 692}{0.526} \approx \frac{300 + 700}{0.5} = 2000$ 'Note that dividing by 0.5 is the same as multiplying by 2'

